

Navigating Wound Balance: Practical approaches for the Australian landscape

Following two recent publications on the concept of Wound Balance (Wounds International, 2023; Blome et al, 2024), an adaptation review was conducted to assess the relevance of Wound Balance in the Australian healthcare context. A Wound Balance Advisory Board was created, consisting of Australian wound care specialists from a range of clinical settings. Two meetings of the Wound Balance Advisory Board took place between 2024 and 2025, with this publication reflecting the outcomes and outlining the journey of acceptance and adaptation of Wound Balance for healthcare professionals (HCPs) in Australia. The inaugural HARTMANN Wound Expert Advisory Board meeting took place in Sydney in August 2024. Participants engaged in pre-reading and surveys designed to explore the potential paradigm shift in mindset and clinical practice introduced by the Wound Balance concept (revolving around 'patient', 'wound' and 'care'). The agenda and objectives for this day-long meeting were clearly defined, ensuring a structured discussion aimed at developing meaningful outcomes.

The second HARTMANN Wound Expert Advisory Board meeting, held in March 2025, focused on refining the insights from the initial meeting to support this publication as an Australian adaptation of Wound Balance. The primary goal was to create a Wound Balance version tailored to include generalist HCPs in Australia and establish consensus on the content. A generalist HCP is defined as 'a HCP who provides wound care as part of their practice, though it is not the primary focus of their daily patient or resident care'. Unlike specialists, generalist HCPs are not exclusively dedicated to wound management.

These discussions explored the importance of refining the terminology and descriptors related to Wound Balance, along with prioritising and promoting 'patient balance' first, not only within the conceptual model, but everyday clinical practice. A key 'take-home' message highlighted the importance of HCP commitment to person-centred, empathetic communication and the development of trusting, therapeutic relationships and compassionate care. Integration of recent international consensus documents and national standards, such as the Therapeutic Wound Cleansing Document: Clinical Evidence and Recommendations (IWII, 2025) and reference to the Australian Standards for Wound Prevention and Management, 4th Edition (2023) was recommended. Opportunities to incorporate Wound Balance as a focused framework for generalist education and resource development was highlighted.

Panel members:

- **Terry Swanson; Co-Chair;** Nurse Practitioner Wound Management, Private Practice; Director and Founder, Wound Education Research Consultancy (W.E.R.C.)
- **Wendy White; Co-Chair;** Independent Consultant/Educator, Advanced Practice Nurse; Director and Founder, TALKING WOUNDS® & WOUNDED® Learning Centre
- **Ann Marie Dunk;** Clinical Nurse Consultant, Tissue Viability Unit, Canberra Health Services
- **Taliesin Ellis;** Wound Management Clinical Practice Consultant, Wound Management Consultancy, Barossa Hills Fleurieu Local Health Network
- **Michelle Gibb;** Nurse Practitioner Wound Management Founder & Director, Wound Specialist Services
- **Chloe Jansz;** Nurse Practitioner/PhD, Director of Nursing Healthcare United, University Lecturer
- **Tabatha Rando;** Nurse Practitioner – Plastics/Wound Management, Plastic and Reconstructive Surgery Department, Royal Adelaide Hospital
- **Sarah Sage;** Clinical Nurse Consultant, Wound Management; Nurse Practitioner, Royal Melbourne Hospital Wound and Lymphoedema CNC, Healthcare United
- **Damian Williams;** Nurse Practitioner, Skin Integrity Services, Royal Brisbane & Women's Hospital

Navigating Wound Balance: Practical approaches for the Australian landscape

What is Wound Balance and why is it important?

What are the barriers to achieving Wound Balance?

The term 'Wound Balance' was originally introduced by a panel of wound management experts in 2023 (Wounds International, 2023), providing a multifactorial wound assessment and management approach that can be tailored in a timely manner to individual needs across different clinical settings. This approach consisted of three pillars, based on the factors that influence wound healing – patient, wound and care [Figure 1]. Wound Balance aims to equip HCPs with an informed approach and practical strategies for managing non-healing wounds that have the potential to heal. However, Wound Balance is equally relevant in addressing wound care imbalances for individuals living with palliative wounds, ensuring tailored and compassionate care.

The Australian group adapted and adopted the following statement from the original white paper introducing the Wound Balance concept (2023): The concept of 'Wound Balance' is multifactorial, encompassing interconnecting considerations related to wound healing and clinical practice. The Wound Balance concept aims to integrate various critical parameters which offer continuity, individualised care and support clinical decision-making, to place

the patient at the centre of all care.' (Wounds International, 2023).

An Australian perspective was proposed by the expert group, expanding the core elements of Wound Balance – the art of person-centric balance, the science of wound and skin balance and the practice of clinical care balance [Table 1]. The Advisory Board members strongly recommend that the message '*Healing starts with me*' must be emphasised. Any HCP responsible for performing wound dressing procedures must recognise that every interaction and intervention carries the potential to impact healing, either positively or negatively. A HCP's attitude, patient communication skills and therapeutic relationships—alongside clinical knowledge and practice—play a crucial role in influencing wound healing outcomes (Probst et al, 2025a).

In Australia, the barriers or challenges for generalist HCPs in planning and delivering optimal wound care via Wound Balance may include the following:

- Lack of confidence
 - This may be related to lack of experience and/or knowledge regarding wound care, wound types, product choices and wound bed preparation
- Lack of understanding of the importance of comprehensive wound care/assessment
 - This may be due to a lack of recognition regarding:
 - Wound management practices as a critical component of undergraduate curricula or as a distinct specialty within healthcare
 - An adequate number of wound care HCPs or specialists to support its integration and advancement
- A lack of awareness or interest in continuously updating knowledge and practice, which may stem from HCPs not recognising that every action—or inaction—can significantly affect the outcomes for individuals living with wounds
- Limited understanding of the science and evidence underpinning Wound Balance
 - Inadequate knowledge of: 1. the essential components of a person-centric wound care approach, resulting in a failure to meet universal healthcare rights




Figure 1

Figure 1. Wound Balance concept (Wounds International, 2023)

Declarations

This advisory board meeting report has been supported by HARTMANN

- and needs for patients; 2. the need to address imbalance of the wound and skin microenvironment; 3. the consequences of delaying implementation of evidence-informed wound care into practice
- Lack of standardised and consistent wound assessment and management practices
 - Not all healthcare facilities in Australia have access to electronic medical records (EMR) and/or standardised wound assessment tools
 - Without an accurate diagnosis and early identification of factors that delay healing, provision of tailored therapy and standard of care (SoC) may be delayed, limited or impossible to achieve
 - Optimal or appropriate treatments may not be available in all care settings
 - Lack of tools for monitoring and documenting patient progress.

More emphasis should be placed on identifying the evidence of delayed or impaired healing and importance of early referrals; to achieve this, it is paramount that HCPs have access to standardised tools for monitoring and documentation. The use of validated QoL or wellbeing Patient Reported Outcomes Measures (PROMS) could be considered by HCPs to better understand the impact of the wound and /or treatment on an individual. To address these challenges, all HCPs should be aware of the factors that may lead to a non-healing wound and how the concept of Wound Balance can help address the barriers to healing.

What prevents wounds from healing?

To understand Wound Balance, it is important for HCPs to understand wound healing physiology and why some wounds transition to non-healing, as well as the impact of imbalance for the patient, wound and care that is delivered and received. A normal wound healing trajectory requires timely occurrence of four consecutive physiological phases: haemostasis, inflammation, proliferation and remodelling (Rodrigues et al 2019; National Library of Medicine [NLM], 2023). However, a large number of patient, wound and care-related factors can impair this normal process, resulting in a non-healing wound that may typically become 'stalled' in the inflammatory stage (Wounds International, 2023; Murray et al, 2018). **Table 2** summarises the factors that impede wound healing.

Although identification of healing barriers, holistic patient and wound assessment and timely, evidence-informed interventions are gold standard approaches for managing non-healing wounds, it can be challenging

Table 1: The core of Wound Balance: The Australian perspective.

Patient	Wound	Care
Person-centric balance	Wound and Skin balance	Clinical Practice balance
Balancing the art: person-centric, empathetic approach to promote healing/optimize symptom control – a standard of care	Balancing the science: micro environments maintained or modulated to promote healing/optimize symptom control – a standard of care	Balancing the practice: evidence-informed screening, assessment, prevention and treatment to promote healing/optimize symptom control – a standard of care

Table 2: Patient, wound and care-related factors that can impede wound healing. Note that several of these factors may be present concomitantly, further complicating the healing progression (adapted from World Union of Wound Healing Societies [WUWHS], 2025).

Person-centric imbalance	Wound and skin imbalance	Clinical practice imbalance
<ul style="list-style-type: none"> • Lack of empathy and compassionate care • Lack of trust in HCPs; compromised therapeutic relationship • Negative emotions (e.g. anger, frustration, depression, anxiety, embarrassment) • Social determinants of health <ul style="list-style-type: none"> - Lack support or engagement/participation with HCP, family friends or significant others • Mis-aligned patient goals/priorities • Negative impact of wound on QoL and wellbeing impacting physical, emotional, social and spiritual/culture • Inadequate health literacy/education <ul style="list-style-type: none"> - Limited disease knowledge • Lack of resources (financial constraints) 	<ul style="list-style-type: none"> • Elevated protease levels (matrix metalloproteinases, polymorphonuclear neutrophil, elastase) • Elevated pro-inflammatory cytokines • Moisture imbalance • Impaired growth factors • Chronic inflammation • M1- vs M2-macrophage imbalance • Biofilm/chronic infection • Cellular burden: (non-viable and/or senescent cells) • Nutrient/oxygen deficiency • Recurrent trauma • Comorbidities/systemic conditions (health status; oxygenation, circulation, perfusion and compromised nutrition) • Elevated pH 	<ul style="list-style-type: none"> • Impact of not recognising contributing health- and age-related factors, and medical conditions that can or have potential to affect healing potential • Limited or inappropriate wound management • Inconsistency in wound care (provider/clinician) • Missed early signs of non-healing • Ritualistic or outdated wound management vs evidence-informed practices • Ineffective education/training • Lack of confidence • Lack of interest; unaware of the impact and influence of appropriate care • Time constraints • Lack of resources

Abbreviations: MMP; matrix metalloproteinases; PMN, polymorphonuclear neutrophils. QoL, quality of life.

in everyday practice to balance all factors contributing to non-healing wounds.

Table 3 provides an explanation of these core components in the Australian wound care context.

Table 3: Explanation of Wound Balance. The first column highlights the Wound Balance pillars and messaging.

Pillars of the Wound Balance concept	Current Components within each pillar	Recommendations for achieving Wound Balance (patient, wound, care)
PATIENT Person-centric Balance Balance the art	Patient-centric assessment and diagnosis	<ul style="list-style-type: none"> • Patient-centric goals may vary and change over time and, therefore, require ongoing empathetic communication, assessment and review of goals on a regular basis • Incorporating culture sensitivity • Encourage engagement and involvement of patient, family and caregivers
	Patient choice	<ul style="list-style-type: none"> • Ensure health literacy assessment and education; demonstrations and review of self-care capabilities for determination of treatment options • Provide patient communication at a level and style/type to meet learning needs • Review understanding and capabilities on an ongoing basis • Cultivate therapeutic relationships and respectful communication that develops trust and engagement
	Balance of patient's QoL	<ul style="list-style-type: none"> • Determine short-term goals based on the immediate need to be controlled such as unmanaged pain or exudate • Explore how the wound and or the treatment has, or is impacting, on their life
	Outcomes measuring – QoL and/or Wellbeing	<ul style="list-style-type: none"> • Consider simple and easy-to-use validated QoL or Wellbeing tool to identify and monitor patient improvement over time (using clinical judgement adhering to local policy/protocols) • Consider cultural-sensitive issues/barriers to healing
WOUND Wound and Skin Balance Balance the science	Early identification and intervention	<ul style="list-style-type: none"> • Identifying the wound type and classification of the wound(s) • Identify factors that increase the risk for chronicity and aetiology on first visit • Standard of Care appropriate for the wound type • Determination of healable, non-healable and maintenance clinical outcomes
	Biomarkers imbalance	<ul style="list-style-type: none"> • Improve knowledge of the signs and symptoms (S&S) of infection using the IWII's Wound Infection Continuum • Assess for S&S of infection during each dressing change • Understand and implement therapeutic wound and skin cleansing • Employ appropriate debridement methods • Ensure appropriate use of antimicrobials (medicated, non-medicated including superabsorbent polymer [SAP]) • Understand the impact of modification and reduction of pH, increased matrix metalloproteases, and wound condition/tissue through dressing selection and wound bed preparation practices
	Manage exudate imbalance	<ul style="list-style-type: none"> • Improve knowledge of the type/consistency and amount of exudate • Improve understanding of chronic wound fluid and the consistency or enzymes within exudate that may contribute to wound and peri-wound damage • Employ appropriate frequency and type of wound dressings for moisture balance
	Normalise (rebalance) and maintain healing trajectory	<ul style="list-style-type: none"> • Employ a standardised and holistic approach to assessment • Monitoring wound healing (percentage reduction in surface area over time); referral to wound specialist/interdisciplinary team when healing does not occur after 30 days/4 weeks • Ensure prevention and management of any recurrent or new wound(s) ce
CARE Clinical Practice Balance Balance the practice	Identify and manage challenges that impact on providing consistent and optimal care (e.g. access to resources)	<ul style="list-style-type: none"> • Develop policies that provide guidance on when to refer a patient and how urgent that review is required • Optimise resources and ensure strong professional networks for collaborative health care • Explore funding options and strategies in providing continuity of care across geographical regions
	Clinical decision and practice continuity of care	<ul style="list-style-type: none"> • Improve knowledge of evidence-informed practice vs habitual management – embrace life-long learning • Review scope of practice for providers and develop early referral patterns when a type of service is not available locally • Identify how to upskill HCPs in the art and science of wound care/management (e.g. compression bandaging and pressure offloading, modification of the non-healing wound microenvironment) • Follow up on diagnostic investigations to promote optimal care
	Realistic time balance in daily practice	<ul style="list-style-type: none"> • During each dressing change, engage in clear discussions to establish specific healing time goals, ensuring alignment with patient needs and clinical expectations • Plan clinical time to allow appropriate time allocation for optimal wound care and interventions

Recommendations for adapting Wound Balance to the Australian context

The following section provides practical recommendations from the panel to assist HCPs align their care goals with each pillar or element of Wound Balance.

Achieving **PATIENT** balance: balancing the art *Person-centric assessment, diagnosis and management to improve engagement*

The goal of improving patient outcomes lies at the heart of Wound Balance. It is, therefore, important to ensure assessment and management is focused on achieving patient-centric balance.

It is important to remember that, despite best practice interventions, not all wounds may heal quickly, and some may never heal (e.g. palliative wounds). Therefore, at every stage of care, using a person-centric, empathetic approach, HCPs and teams can commit to optimising patient comfort, developing and maintaining therapeutic relationships through effective communication and maintaining patient partnerships and engagement (Probst et al, 2025a; Probst et al, 2025b). The Australian Advisory Board proposed the Wound Balance mantra, *'Healing starts with me'*.

Measurement and monitoring of QoL or wellbeing can be considered an additional 'tool in the tool shed' for person-centric assessment and management interventions (Janke et al, 2023; Upton et al, 2021). The use of validated wound-specific tools is guided by local policy or protocols and HCP or team judgement, taking into consideration available resources along with patient and setting suitability.

Focus on patient goals

Patient goals may vary significantly over time; it is important that assessment and reviews are undertaken routinely.

- It is important to incorporate cultural sensitivity in all patient interactions and communications. A significant number of patient communications may occur as 'bedside conversations', for which, HCPs should receive appropriate training and education
- There is a need to provide appropriate patient education materials (e.g. health literacy and cultural requirements)
- It is crucial to foster referral pathways for identified patient-related issues, including physical, social and emotional spiritual/cultural needs.

Achieving **WOUND** and **SKIN** balance: balancing the science

Wound pathology and biomarkers


For achieving wound care balance, it is crucial to understand the shift in wound pathology (as indicated by both biomarkers and clinical signs and symptoms) at each stage of wound healing. Furthermore, due to the long-term nature of a non-healing wound, it is imperative that HCPs are able to recognise wound infection and chronic inflammation and understand their local referral pathway as well as guidelines for antibiotic prescription (for localised, spreading and/or systemic infections) and antimicrobial stewardship strategies. The panel recommends that HCPs employ the Wound Infection Continuum for identifying signs and symptoms of wound infection (IWII, 2022) and identification and management of non-viable tissue; it is important that HCPs understand the concept of debridement and the importance of an aseptic approach to wound care procedures.

Early identification and intervention

Upon presentation, it is important to identify the wound type and classification, and assess the risk factors related to patient, wound and care imbalance involved in wound chronicity. Recognising healing barriers is essential, as their early identification can help detect wounds at risk of becoming non-healing. Accurate recognition of these factors enables HCPs to identify potential red flags and ensures timely intervention or referral. In Australian care settings, there is a clear need for a standardised approach to wound prevention, assessment, and management, as outlined in the Australian Standards for Wound Prevention and Management (2023).

Based on Wound Balance, consideration of the following factors and recommendations can help HCPs develop a standardised strategy toward early identification and suitable interventions for non-healing wounds:

1. Holistic assessment and identifying factors that may delay healing (physical, emotional, social and spiritual/cultural)
 - Patient factors
 - Comorbidities
 - Psychosocial
 - QoL or Wellbeing
2. Barriers in identification of wound type and or classification
 - If wound type and/or classification is unclear, HCPs should seek specialist advice aimed at further diagnostic interventions and assessment
 - If the practice or service is unable to identify and classify the wound type, an



Scan the QR code to access the IWII publication and view the Wound Infection Continuum

appropriate referral should be made to a wound care specialist, or appropriate team member(s)

3. Identification of 'red flags' that require referral
 - A referral may be 'emergency', that is, emergency intervention/care is required due to medical need and life- or limb-threatening
 - Alternately, an 'urgent' referral may be needed, due to unmanaged or escalating needs such as pain, exposed structures or decreased vascularity
 - A referral may be 'non-urgent' where further diagnostic assessments are required to ensure tailored/targeted therapy
4. Targeted treatment based on wound type and/or classification
 - Wound bed preparation (macro and micro wound environment) and use of validated wound assessment tools, such as TIMERS (Wounds International, 2023)
 - Appropriate skin products, dressings or devices for clinical need and patient preferences
 - SoC, based on wound type and patient preference
5. Monitoring of healing trajectory and re-assessment on a regular basis
 - If the wound is improving, care should be continued
 - If the wound is not healing, re-assess for factors delaying healing, or refer to specialist services.

Normalising wound healing trajectory and balancing exudate

The goal of Wound Balance is to minimise factors that hinder wound healing while enhancing those that support recovery in healable wounds. Recognising expected healing trajectories for various wound types is essential—when healing is slow or absent, an investigation into the underlying causes is necessary. In cases of non-healing wounds with healing potential, HCPs must take proactive measures to modify the microenvironment and optimise conditions for recovery. This approach aids clinicians in effectively managing their time, ensuring appropriate allocation for wound dressing procedures, and implementing proactive treatment strategies.

An understanding of healing biomarkers can assist in placing a non-healing wound on a healing trajectory. Recent studies have demonstrated a significant difference in the expression of certain protein biomarkers between healing and non-healing wounds

(Stacey et al, 2019; Mikosiński et al, 2022). In a group of patients with highly exuding wounds (venous leg ulcers; $n=57$), nine healing biomarkers were found to be significantly different between healing versus non-healing wounds (Mikosiński et al, 2022). Furthermore, Mikosiński et al (2022) demonstrated that biomarker profiles may be useful in early identification of patients at risk of developing non-healing wounds.

It is important to assess the type and consistency of exudate, as well as understand the proteolytic enzymes within it (e.g. metalloproteases) that may compromise the integrity of peri-wound skin and damage growth factors and wound substrate. To achieve Wound Balance, HCPs may consider easy-to-use, effective, first-response dressings that can modify the imbalanced microenvironment, balance wound exudate, minimise wound-related pain while reducing skin injury risk via atraumatic removal. Wound dressings containing superabsorbent polymers (SAP) with these characteristics are now available; these dressings can assist in moving the wound toward a healing trajectory (Wounds International, 2023). SAP dressings maintain a moist wound environment, offer high absorbance capacity and reduce wound healing inhibitors, such as matrix metalloproteinase 2 and elastase, and microorganisms (Probst, 2019; Candas et al, 2021; Veličković et al, 2024). The moist wound surface promotes autolytic debridement, further removing dead tissue and debris (Ousey et al, 2016; Choo et al, 2019).

Therefore, SAP dressings can help achieve Wound Balance in most clinical settings by promoting a shift to healing (Wound Balance Wounds International, 2023; WUWHS, 2025).

Achieving CARE balance: balancing the practice Addressing clinical practice-related challenges

Referral and escalation pathways may vary significantly across different healthcare settings. It is important to ensure that appropriate referral and escalation pathways are in place and HCPs are informed.

The Advisory Board recommends nurturing strong professional networks between generalist HCPs and experts who undertake wound care. This collaborative environment can help optimise resources and promote timely interventions and, therefore, improve patient outcomes.

There is also a need to ensure HCPs can define and differentiate between 'urgent', 'new' and 'non-urgent' referral needs.

Clinical decision-making and practicing continuity of care

Given that non-healing wounds are a long-term condition, it is important to ensure continuity of care using an evidence-informed approach. All generalist HCPs undertaking wound care should be aware of their local referral pathways and wound prevention and management standards and guidelines (e.g. for adjunct therapies/devices/diagnostics). Continuity of care is best achieved through a multidisciplinary approach that includes allied health professionals such as podiatrists, dietitians and physiotherapists. Collaborative teamwork enhances patient outcomes by integrating expertise and holistic care strategies.

Further recommendations were provided by the Advisory Board to highlight the variety of funding/support available for ensuring wound care continuity. Australian government or non-government funding schemes or support services, can be accessed by eligible populations living in the primary health care setting. This may include (but is not limited to):

- My Aged Care services
- Home and Community Care (HCCA), National Disability Insurance Scheme (NDIS), Department of Veterans Affairs (DVA) and Repatriation Pharmaceutical Benefits Scheme (RPBS) or Chronic Wound Consumables Scheme (CWSC 2025).

Realistic time balance in daily practice

The panel recommends achieving time balance by undertaking clear discussions among generalist HCPs to establish specific care goals and expected healing timelines for each patient. These goals should be re-assessed and concerns addressed at each dressing change.

To ensure continuity and timely communication between HCPs (including local and remote team members), thorough clinical documentation can play a crucial role. For example, assessment at each dressing change can be documented, recording wound and skin assessment findings including measurements and photographs to monitor healing progression. Early identification of red flags can trigger wound specialist referral if the wound does not heal in four weeks (Atkin et al, 2019).

The significant gaps in the Australian wound care landscape can be addressed by implementing Wound Balance

Non-healing wounds remain a significant healthcare challenge across the world (AMA, 2022; Sen, 2023; Sharma et al, 2024) and are referred to as a 'silent epidemic' by the Australian federal government, peak bodies

and professional organisations. Carville et al (2022) found that, with no added costs to the payers, healing for up to 80% of all non-healing wounds may be achieved with implementation of evidence-informed care via contemporary wound care products. However, the burden of care for non-healing wounds is still high, costing Australia approximately \$5.14 billion in 2019 (Queen and Harding, 2023).

The Australian Health Research Alliance (AHRA) underlines the importance of equitable wound care for all patients, regardless of 'their geographic location, type of wound or healthcare provider (AHRA, 2025). Several Australia-specific cultural and demographic barriers currently exist in providing inclusive and equitable wound care. A recent consensus highlighted the national need to focus research on non-healing wounds, with particular emphasis on managing pain, venous leg ulcers and pressure injuries (Finlayson et al, 2025). Smith et al (2024) demonstrated that integrating wound care data across hospitals and community settings in Australia enhances HCPs' understanding of associated comorbidities and the diverse outcomes across different wound aetiologies.

Data from older and more vulnerable populations indicate that, in Australia, concurrent use of complementary medicines with prescription treatments has not changed in the last few years and presents a significant safety concern (Harnett et al, 2023).

Furthermore, for certain patient groups (e.g. First Nation populations), clinicians may require assistance from or collaboration with traditional healers. The impact of rural/semi-rural living on access to wound care facilities is also significant.

These data highlight an unmet need in Australia to implement evidence-informed wound care practices, tailored to addressing the local needs (McCosker et al, 2018; Pacella et al, 2022). In summary, these barriers impact patients' access to evidence-based wound care, likely leading to sub-optimal outcomes.

The concept of Wound Balance provides a holistic, tailorable approach that generalist HCPs in Australia require for improving healing outcomes for their patients and reducing long-term wound care costs.

Conclusions and future recommendations

This publication highlights the role that Wound Balance can play in addressing the unmet wound care needs in Australia by empowering both the generalist HCPs and



Scan the QR code above to access the global consensus publication on Wound Balance, 'Implementing Wound Balance: Outcomes and future recommendations' (WUWHS, 2025)

specialists in wound care. Furthermore, the recommendations from the Advisory Board on each pillar of Wound Balance provide a foundation upon which future wound care education programmes can be developed for generalist HCPs and a sustainable improvement achieved.

With an ageing population and the projected rise in the prevalence of non-healing wounds in Australia (AHRA, 2025), there is a significant need to educate HCPs about the Wound Balance concept for prompt implementation of evidence-informed wound care strategies. Wound Balance can help create a roadmap for Australian educators and HCPs in providing equitable, evidence-informed care—with patient, wound and care balance for all.

Further reading: The 2nd Australian Advisory Board met on 7th March 2025 and, since this meeting, a global consensus on Wound Balance has been published (WUWHS, 2025). Scan the QR code on this page to access the consensus publication.

References

- Australian Health Research Alliance (2025) Wound care. Available at: <https://ahra.org.au/our-work/wound-care/> (accessed 24.04.2025)
- Australian Medical Association (2022) Solutions to the chronic wound problem in Australia. Available at: <https://www.ama.com.au/sites/default/files/2022-10/Solutions%20to%20the%20chronic%20wound%20problem%20in%20Australia.pdf> (accessed 22.04.2025)
- Atkin L, Bućko Z, Conde Montero E et al (2019) Implementing TIMERS: the race against hard-to-heal wounds. *J Wound Care* 28(3 Suppl 3): S1–49
- Australian Standards for Wound Prevention and Management, 4th edition (2023) Available at: <https://woundsaustralia.org/int/woundsaus/uploads/Publications/Standards%20and%20Guidelines/Australian%20Standards%20for%20Wound%20Prevention%20and%20Management%204th%202023.pdf> (accessed 16.05.2025)
- Blome C, Candas E, Erfert-Berge C et al (2024) Implementing the Wound Balance concept into routine practice worldwide. *Wounds Int* 15(2): 44–50
- Candas E, Forster J, Schäfer J et al (2021) HydroClean® Made Easy. *Wounds International*
- Choo J, Nixon J, Nelson A (2019) Autolytic debridement for pressure ulcers. *Cochrane Database Syst Rev* 2019(6): CD011331
- Finlayson K, Haesler E, Bui UT et al (2025) Priority topics for chronic wound research in Australia: a consensus study. *Wound Prac Res* 33(1): 5–17
- Harnett J, McIntyre E, Adams J et al (2023) Prevalence and Characteristics of Australians Complementary Medicine Product Use, and Concurrent Use with Prescription and Over-the-Counter Medications—A Cross Sectional Study. *Nutrients* 15(2): 327
- International Wound Infection Institute (2022) Wound Infection in Clinical Practice. *Wounds International*
- International Wound Infection Institute (2025) Therapeutic wound and skin cleansing: Clinical evidence and recommendations. *Wounds International*
- Janke TM, Kozon V, Valiukeviciene S et al (2023) Validation of the Wound-QoL-17 and the Wound-QoL-14 in a European sample of 305 patients with chronic wounds. *Int Wound J* 21(3): e14505
- McCosker L, Tulleners R, Cheng Q et al (2019) Chronic wounds in Australia: A systematic review of key epidemiological and clinical parameters. *Int Wound J* 16(1): 84–95
- Mikosiński J, Kalogeropoulos K, Bundgaard L (2022) Longitudinal Evaluation of Biomarkers in Wound Fluids from Venous Leg Ulcers and Split-thickness Skin Graft Donor Site Wounds Treated with a Protease-modulating Wound Dressing. *Acta Derm Venereol* 102: adv00834
- Murray RZ, West ZE, McGuinness W (2018) The multifactorial formation of chronic wounds. *Wound Practice and Research* 26(1): 38–6
- National Library of Medicine (2023) Wound Healing Phases. Available at: https://www.ncbi.nlm.nih.gov/books/NBK470443/#_NBK470443_dtlis__ (accessed 22.04.2025)
- Ousey K, Rogers AA, Rippon MG (2016) HydroClean® plus: a new perspective to wound cleansing and debridement. *Wounds UK* 12(1): 78–87
- Pacella RE, Tulleners R, Cheng Q et al (2018) Solutions to the chronic wounds problem in Australia: a call to action. *Wound Prac Res* 26(2): 84–98
- Probst A (2019) Zetuvit Plus Silicone Border Made Easy. *Wounds International*
- Probst S, Menon T, Stefanelli A et al (2025a) Empathy in Wound Care: A Scoping Review of Its Role, Impact, and Barriers to Person-Centred Healing. *Int Wound J* 22: e70687
- Probst S, Bergin SM, Tehan P (2025b) Healing beyond the surface: Empathy's role in chronic wound care a qualitative study. *J Tissue Viability* 34(3): 100910
- Queen D, Harding K (2023) What's the true costs of wounds faced by different health systems around the world? *Int Wound J* 20: 3935–8
- Rodrigues M, Kosaric N, Bonham CA et al (2019) Wound Healing: A Cellular Perspective. *Physiol Rev* 99(1): 665–706
- Sen CK (2023) Human Wound and Its Burden: Updated 2022 Compendium of Estimates. *Adv Wound Care* 12: 657–70
- Sharma A, Shankar R, Yadav AK et al (2024) Burden of Chronic Nonhealing Wounds: An Overview of the Worldwide Humanistic and Economic Burden to the Healthcare System. *Int J Low Extrem Wounds* 24: 15347346241246339
- Smith J, Carville K, Smith K et al (2024) The impact of comorbidities on wound healing in the community: the value of using linked hospital data. *Wound Prac Res* 32(3): 136–44
- Stacey MC, Phillips SA, Farrokhhyar F (2019) Evaluation of wound fluid biomarkers to determine healing in adults with venous leg ulcers: a prospective study. *Wound Repair Regen* 27: 509–18
- Trouth S (2024) Introducing Wound Balance: placing the patient at the heart of wound healing. *Wounds UK* 20(1): 32–7
- Upton P, Cartwright M, Upton D (2021) Living with chronic wounds: an exploration of adaptive and maladaptive coping strategies and their association with wellbeing. *Wounds International* 12(1): 12–7
- Veličković V M, Macmillan T, Lones E (2024) Systematic review and quality assessment of clinical and economic evidence for superabsorbent wound dressings in a population with chronic ulcers. *Int Wound J* 21(3): e14750
- Wounds International (2023) *Wound Balance: achieving wound healing with confidence*. Wounds International
- World Union of Wound Healing Societies (2025) *Implementing Wound Balance: Outcomes and future recommendations*. Wounds International