

The role of wound cleansing in the management of wounds



Authors:
Randall Wolcott, Jacqui Fletcher

Wound cleansing should be seen as an integral part of wound bed preparation to optimise the wound environment by removing debris, reducing bacterial load and preventing biofilm activity. Clinicians have a number of options to choose from when selecting an appropriate wound irrigation solution. One factor to consider includes the ability to mount a rapid antimicrobial response to microbial contamination, while avoiding damage to human cells important for wound healing.

Micro-organisms produce significant barriers to the healing of chronic wounds^[1]. They are present in all wounds, although not all wounds become chronic. Microbes grow in both a planktonic phenotype (free-floating mobile single cells) and biofilm phenotype (a fixed polymicrobial community)^[2]. A self-secreted matrix protects the biofilm from attack by the host immune system and makes the wound harder to treat^[3]. In fact, many of the characteristics of chronic wounds, such as persistent inflammation, exudate, and host-cell senescence, result directly from biofilm^[4,5,6].

Wounds that have become chronic — characterised by delayed or stalled healing — due to suspected biofilm/increased bioburden should be managed with a biofilm-based approach to wound care^[7]. This means combining wound cleansing/irrigation with debridement and application of topical antimicrobial agents (e.g. wound cleansers and dressings), to facilitate healing by disrupting, removing and preventing the reformation of biofilm^[8].

Importance of wound bed preparation

Wound bed preparation is recognised as having a key role in wound management. The International Advisory Board on Wound Bed Preparation has developed an assessment tool, known by the acronym TIME (T = tissue, non-viable or deficient; I = infection or inflammation; M = moisture imbalance; E = edge of wound, non-advancing or undermined), that sets out the goals of wound-bed preparation: removing non-viable tissue, reducing oedema and exudate, reducing the bacterial burden and correcting any abnormalities to promote wound healing^[9]. Wound bed preparation offers a

structured and systematic approach to assist clinicians when assessing and managing patients with wounds.

Where wound cleansing fits

Wound cleansing [Box 1] can help achieve the goals of wound bed preparation by assisting in removing loose material to create the optimal local conditions for wound healing by removing exudate and other debris. Irrigation is the preferred method of wound cleansing, as it can clear the wound of debris and microbes, while avoiding trauma in the wound bed^[10].

A practical strategy for wound bed preparation in chronic wounds is to lightly irrigate the wound before inspection and assessment. Debride any soft, degraded areas of the wound bed, slough or necrotic tissue according to local protocols, and then re-irrigate the wound with a cleansing solution before applying an appropriate dressing, also according to local protocols.

There are several broad categories of solutions that can be used: saline and water, highly reactive solutions, and minimally or non-cytotoxic antimicrobial-containing solutions.

Exploring irrigation solutions

When considering the clinical benefits of wound irrigation and the appropriate agents to use, the clinician should keep in mind that an ideal wound irrigation solution will provide periodic reduction of bacterial contamination and removal of debris without adversely affecting cellular activities crucial to the wound-healing process^[11]. It is, therefore, important to consider the balance of antimicrobial action and cytotoxicity when choosing a wound irrigation solution [Table 1].

Saline and tap water

Although tap water and saline are not

Randall D Wolcott is Medical Director, Southwest Regional Wound Care Center, Lubbock, USA and Jacqui Fletcher is Clinical Strategy Director, Welsh Wound Innovation Centre, Cardiff, UK

IT'S TIME FOR A

REVOLUTION

IT'S TIME FOR WOUNDOX® IRRIGATION SOLUTION



Woundox® Irrigation Solution contains hypochlorous acid, a powerful antimicrobial agent that:

- > Rapidly facilitates the reduction of bacterial bioburden^{1,2,3}
- > Has wide efficacy against microbes that inhibit wound healing^{1,2}
- > Helps to prevent the creation of resistant strains by attacking bacteria at multiple sites⁴
- > Is non-irritating, non-sensitising, non-cytotoxic and does not negatively impact the wound healing process^{3,5}
- > Facilitates the reduction of harmful microbes on the wound surface^{1,2,3}

Woundox® Irrigation Solution's active ingredient promotes healthy tissue⁶

Become part of the solution
visit woundcarerevolution.co.uk

References:

1. Clinical evaluation report. Data on file. martindale Pharma.
2. Shetty N et al. Journal of Hospital Infection 1999; 41: 101-105.
3. Niezgodka J et al. Adv Skin Wound Care 2010; 23: 352-357
4. Zinkevich V et al. Journal of Hospital Infection 2000; 46: 153-156.
5. Biological Evaluation Report. Data on File, Martindale Pharma.
6. Wang L et al. J Burns Wounds 2007; 6: 65-79

wcycytotoxic and do not seem to be harmful to wounds^[12], these cleansing choices may not actively promote healing, particularly in chronic wounds with biofilm and/or increased bacterial burden^[13], indicating that non-antiseptic cleansing does not remove harmful molecules such as matrix metalloproteases and elevated proinflammatory cytokines. These molecules may be present in chronic wounds and delay healing.

Although these agents have been used as standards of non-cytotoxicity^[14], neither has significant antimicrobial properties. This means saline is not appropriate for augmenting wound-bed preparation in the context of biofilm-based wound care^[15], or where wounds are clinically diagnosed as locally infected or at risk of infection. Further, it should be pointed out that even 'safe' tap water can become colonised with viable microbes. In particular, *Pseudomonas* is well-documented in the plumbing systems of healthcare facilities^[16,17,18]. Therefore, a cleansing agent with broad-spectrum activity and rapid kill rates (dwell time of 5 to 10 minutes) should be chosen to aid wound bed preparation.

Highly reactive solutions

Solutions such as peroxide and povidone iodine (depending on the carrier and the strength of the povidone iodine) and commercially available products (e.g. alcohol-based cleansers, soaps, foams and wipes) can be particularly cytotoxic^[19]. A 2010 Cochrane review suggested that such solutions may do little to control wound bacteria and may in fact interfere with host healing mechanisms^[12]. Although they probably do not significantly harm the host, the high level of reactivity is not necessary.

Antimicrobial solutions

Multiple antiseptic agents are available and have been extensively evaluated for cytotoxicity and their biocidal abilities against a broad spectrum of microorganisms, including bacteria and fungus (yeast)^[15]. Common cleansing agents such as octenidine

and polyhexamethylene biguanide (PHMB) are minimally toxic to host cells when used in low concentrations^[21].

Hypochlorous acid is produced by the body's immune cells in response to invading pathogens^[22]. It is available as a commercially-prepared wound cleansing solution, which is non-toxic, effective against a broad range of microorganisms and has a rapid kill rate^[23,24]. Clinical studies of hypochlorous acid demonstrate improved wound-healing outcomes^[24,25].

Rationale for antimicrobial solutions

After a wound has been washed and debrided, exposing microbial cells to the environment, the antimicrobial properties of a cleansing agent can be effective. Because microbes may exist on the wound bed in high numbers, agents stronger than simple saline or tap water may be needed. Antimicrobial solutions can be used as part of the balance in managing wound colonisation, biofilm and/or infection. Non-toxic agents that are effective at low concentrations should be considered when possible.

Antimicrobial solutions have the added benefit of being usable across the spectrum of chronic wounds without resulting in bacterial resistance issues, because they are antiseptic rather than antibiotic. These factors, along with the ability to disrupt and prevent reformation of biofilm/reduce bioburden are ideal properties of an antimicrobial irrigation solution [Figure 1]. As a result, such irrigation solutions may have an important place within the process of chronic wound management.

Summary

Resolving critical colonisation, biofilm and/or infection in a chronic wound is achieved through appropriate wound bed preparation. The goal of wound cleansing is to remove loose material in the wound bed, reduce bacterial load and to assist in the suppression of biofilm.

Cleansing should be tailored according to the goals determined by holistic assessment of the patient and wound, and best practices for irrigation followed [Box 2]. Generally speaking, wounds should be irrigated at every dressing change until visible debris is removed. Cleansing agents may also be useful in removing encrusted dressings in order to lift them and avoid causing trauma to newly formed granulation tissue in the wound bed.

It is critical that the wound cleansing agent does not impair the wound healing process,

Box 1. Definition of wound cleansing

- Remove surface contaminants, loose debris, slough, softened necrosis, microbes and/or remnants of previous dressings from the wound surface and its surrounding skin^[20].

Table 1. Key considerations in some common wound irrigation agents.

Saline ^[10]	<p>Low toxicity</p> <p>Limited ability to reduce bacterial load</p> <p>Bacterial growth can occur in an open container within 24 hours</p>
Sterile water ^[10]	<p>Limited ability to reduce bacterial load</p> <p>Readily absorbed by tissues; water toxicity may result when excess volumes are used</p> <p>No longer sterile after opening</p>
Tap water ^[10,17]	<p>Recommended where saline and sterile water are not available</p> <p>Limited ability to reduce bacterial load</p> <p>Microbes, in particular <i>P. aeruginosa</i>, can colonise taps and as a result may end up in wounds irrigated in this way</p>
Commercially available products (e.g. foams, soaps, wipes and solutions with surfactants) ^[10]	<p>Remove bacteria with less required force due to surfactant content</p> <p>May be best suited for wounds with adherent cellular debris and biofilm</p> <p>Typically contain preservatives to extend effective shelf life</p> <p>Can be highly cytotoxic to healthy cells and granulating tissue</p>
Povidone iodine ^[10]	<p>Broad-spectrum antimicrobial activity</p> <p>Cytotoxic to healthy cells and granulating tissue in higher-percentage concentrations</p> <p>May irritate periwound skin</p>
Hydrogen peroxide ^[10]	<p>May be cytotoxic to healthy cells and granulating tissue</p> <p>Ineffective in reducing bacterial counts</p>
Polyhexamethylene biguanide (PHMB) 0.1% ^[26]	<p>Also contains betaine, a surfactant, to lift microbes and debris and suspend them in solution to prevent wound recontamination</p> <p>Has an increased ability to penetrate difficult-to-remove coatings, lifting debris, bacteria and biofilm from the wound</p> <p>Broad spectrum of activity against bacteria, viruses and fungi</p> <p>No evidence of resistance</p>
Octenidine ^[27]	<p>Can be used to loosen encrusted dressings in addition to irrigating debris and microbes from the wound bed</p> <p>Contains octenidine dihydrochloride, a preservative, to extend shelf life and a surfactant-like molecule that is effective at infiltrating wounds while being less irritating</p> <p>Shown to prevent and remove the growth of bacterial biofilms</p>
Hypochlorous acid 0.01% ^[11,23,28,29,30,32]	<p>Broad-spectrum antimicrobial activity</p> <p>Non-irritating, non-sensitising, non-toxic</p> <p>Can be used to loosen encrusted dressings in addition to irrigating loose debris and bacteria from the wound bed</p> <p>Has rapid antimicrobial activity at concentrations safe for human cells</p>

but is strong enough to remove the protective matrix secreted by microbes attached to the wound (biofilm) and reduce the bioburden.

Clinicians have a number of options when selecting a wound cleansing agent [Figure 1]. Although saline and water have been found by numerous studies to not be harmful to wounds, this feature is not enough in the context of biofilm-based wound care. Other wound cleansers, such as povidone iodine (in higher concentrations), soaps, peroxide and alcohol may be too harsh (Rabenberg et al, 2002)^[33].

The ideal wound irrigation solution should exhibit potent and rapid antimicrobial activity at concentrations that do not damage host cells required for wound healing. Hypochlorous acid,

when used in low concentrations, is non-toxic and has a rapid antimicrobial action. The choice of irrigation solution should also reflect the individual requirements of the wound and the patient after thorough holistic assessment^[13].

Where holistic assessment leads to the suspicion that bioburden is delaying wound healing, clinicians should consider using a suitable antimicrobial wound irrigation solution to promote healing, reduce bioburden, prevent the proliferation of biofilms and/or remove biofilm before applying an appropriate dressing. As such, cleansing should become an integral part of wound-bed preparation, to help prevent infection and optimise the wound environment for healing.

WINT

References

- Kim M, Ashida H, Ogawa M. Bacterial interactions with the host epithelium. *Cell Host Microbe* 2010; 8(1): 20–35
- James GA, Swogger E, Wolcott R et al. Biofilms in chronic wounds. *Wound Repair Regen* 2008; 16(1): 37–44
- Tuttle MS, Mostow E, Mukherjee P. Characterization of bacterial communities in venous insufficiency wounds by use of conventional culture and molecular diagnostic methods. *J Clin Microbiol* 2011; 49(11): 3812–9
- Katsuyama M, Ichikawa H, Ogawa S, Ikezawa Z. A novel method to control the balance of skin microflora. Part 1. Attack on biofilm of *Staphylococcus aureus* without antibiotics. *J Dermatol Sci* 2005; 38(3): 197–205
- Wolcott RD, Rhoads DD, Dowd SE. Biofilms and chronic wound inflammation. *J Wound Care* 2008; 17(8): 333–41
- Raymond B, Young JC, Pallett M et al. Subversion of trafficking, apoptosis, and innate immunity by type III secretion system effectors. *Trends Microbiol* 2013; 21(8): 430–41
- Wolcott RD, Rhoads DD, Bennett ME et al. Chronic wounds and the medical biofilm paradigm. *J Wound Care* 2010; 19(2): 45–53
- Rosdahl C, Kowalski M. *Textbook of Basic Nursing*. (9th edn). Lippincott, Williams & Wilkins, 2008: 769
- Falanga V. Wound bed preparation: science applied




Figure 1. Wound-cleansing decision making pathway.

Box 2. Practical tips for wound irrigation.

- Choice of solution: Base choice on assessment of both the patient (including medical condition and allergies) and the wound (e.g. clinically assessed as critically colonised^[10,34]
- Method of delivery: Deliver irrigation based on the needs of the patient (e.g. pain levels) and wound (e.g. fragility of wound and periwound skin^[10,34]).
- Volume of solution: Volumes of 50–100 ml per centimetre of wound length is the general rule of thumb^[35,36].
- Prevention of cross-contamination: The clinician should wear personal protective equipment. Do not use solution that has been opened for longer than 24 hours^[10,34].
- Comfort of patient: Make sure irrigation solution is at room temperature or slightly warmer. Use analgesia for painful wounds and allow time for it to take effect^[10,34].
- Irrigation of wound: Position the patient so the solution runs from the upper end of the wound downward or from clean to dirty (if the upper end is heavily infected and the lower end is clean), into a clean basin or irrigating pouch^[10,34].
- Documentation of treatment: Record all aspects of the wound cleansing, including assessment of the wound (e.g. slough, exudate, pain, erythema), date and time of treatment, amount and type of solution used, skin care performed, wound dressing(s) applied and notes on the patient's concordance with treatment^[10,34].

to practice. In: *European Wound Management Association position document: Wound bed preparation in practice*. MEP Ltd, 2004

10. Gabriel A, Schraga ED, Windle ML (2013) *Wound irrigation*. Medscape. <http://emedicine.medscape.com/article/1895071-overview> (accessed 10.06.2014)
11. Hoon R, Rani SA, Najafi R et al. *Antimicrobial activity comparison of pure hypochlorous acid (0.01%) with other wound and skin cleansers at non-toxic concentrations*. Poster presented at: Symposium on Advanced Wound Care Spring and Wound Healing Society, Denver, USA, 1–5 May 2013
12. Fernandez R, Griffiths R, Ussia C. Water for wound cleansing. *Cochrane Database Syst Rev* 2002; (4): CD003861
13. Cutting K. Addressing the challenge of wound cleansing in the modern era. *Br J Nurs* 2010; 9(11): S24–9
14. Wilson JR, Mills JG, Prather ID, Dimitrijevic SD. A toxicity index of skin and wound cleansers used on in vitro fibroblasts and keratinocytes. *Adv Skin Wound Care* 2005; 18(7): 373–8
15. Leaper DJ, Schultz G, Carville K, et al. Extending the TIME concept: what have we learned in the past 10 years? *Int Wound J* 2012; 9(Suppl 2): 1–19
16. Trautmann M, Lepper PM, Haller M. Ecology of *Pseudomonas aeruginosa* in the intensive care unit and the evolving role of water outlets as a reservoir of the organism. *Am J Infect Control* 2005; 33(5 Suppl 1): S41–9
17. Mena KD, Gerba CP. Risk assessment of *Pseudomonas aeruginosa* in water. *Rev Environ Contam Toxicol* 2009; 201: 71–115
18. Jefferies JM, Cooper T, Yam T, Clarke SC. *Pseudomonas aeruginosa* outbreaks in the neonatal intensive care unit – a systematic review of risk factors and environmental sources. *J Med Microbiol* 2012; 61(Pt 8): 1052–61
19. Sibbald RG, Leaper DJ, Queen D (2011) Iodine made easy. *Wounds International* 2011; 2(2). Available at: <http://www.woundsinternational.com/made-easys/iodine-made-easy> (accessed 24.09.2014)
20. Rodeheaver RT, Ratliff CR. Wound cleansing, wound irrigation and wound disinfection. In: Rodeheaver GT, Krasner DL, Sibbald RG eds. *Chronic Wound Care: A Clinical Source Book for Healthcare Professionals*. HMP Communications, Malvern, USA, 2007
21. Rani SA, Hoon R, Najafi RR, et al. The in vitro antimicrobial activity of wound and skin cleansers at nontoxic concentrations. *Adv Skin Wound Care* 2014; 27(2): 65–9
22. Eryilmaz M, Palabruyik IM. Hypochlorous acid – analytical methods and antimicrobial activity. *Topical J Pharm Res* 2013; 12(1): 123–6
23. Wang L, Bassiri M, Najafi R. Hypochlorous acid as a potential wound care agent: Part 1. Stabilized hypochlorous acid: a component of the inorganic armamentarium of innate immunity. *J Burns Wounds* 2007; 6: e5
24. Crew J, Varilla R, Rocas TA et al. NeutroPhase® in chronic non-healing wounds. *Int J Burns Trauma* 2012; 2(3): 126–34
25. Robson MC, Payne WG, Ko F. Hypochlorous acid as a potential wound care agent: Part II. Stabilized hypochlorous acid: Its role in decreasing tissue bacterial bioburden and overcoming the inhibition of infection on wound healing. *J Burns Wounds* 2007; 6: e6
26. Bradbury S, Fletcher J. Prontosan made easy. *Wounds International* 2011; 2(2). Available at: www.woundsinternational.com/made-easys/prontosan-made-easy (accessed 24.09.2014)
27. Braun M, McGrath A, Downie F (2013) Octenilin range made easy. *Wounds UK* 2013; 9(4): 1–4. Available at: www.wounds-uk.com/made-easy (accessed 24.09.2014)
28. Shetty N, Srinivasan S, Holton J, Ridgway GL. Evaluation of microbicidal activity of a new disinfectant: Sterilox 2500 against *Clostridium difficile* spores, *Helicobacter pylori*, vancomycin resistant *Enterococcus* species, *Candida albicans* and several *Mycobacterium* species. *J Hosp Infect* 1999; 41(2): 101–5
29. Zinkevich V, Beech IB, Tapper R, Bogdarina I. The effect of super-oxidized water on *Escherichia coli*. *J Hosp Infect* 2000; 46(2): 153–6
30. Selkon JB, Cherry GW, Wilson GM, Hughes MA. Evaluation of hypochlorous acid washes in the treatment of chronic venous leg ulcers. *J Wound Care* 2006; 15(1): 33–7
31. Niezgoda JA, Sordi PJ, Hermans MH (2010) Evaluation of Vashe Wound Therapy in the clinical management of patients with chronic wounds. *Adv Skin Wound Care* 23(8): 352–7
32. Martindale Pharma. Clinical evaluation report. Data on file.
33. Rabenberg VS, Ingersoll CD, Sandrey MA, Johnson MT (2002) The bactericidal and cytotoxic effects of antimicrobial wound cleansers. *J Athl Train* 37(1): 51–4
34. Sardina D. Best practices: Is your wound-cleansing practice up to date? *Wound Care Advisor* 2013; 2(3): 15–7. Available at http://woundcareadvisor.com/is-your-wound-cleansing-practice-up-to-date_vol2_no3 (accessed 10.06.2014)
35. Chisholm CD, Cordell WH, Rogers K, Woods JR. Comparison of a new pressurized saline canister versus syringe irrigation for laceration cleansing in the emergency department. *Ann Emerg Med* 1992; 21(11): 1364–7
36. Lammers RL, Hudson DL, Seaman ME. Prediction of traumatic wound infection with a neural network-derived decision model. *Am J Emerg Med* 2003; 21(1): 1–7

Acknowledgement

This article has been supported by Martindale Pharma.

Expert commentary: wound cleansing

The microbiological profile of chronic wounds is diverse, and Wolcott and Fletcher highlight the developing evidence base regarding the role of planktonic and biofilm phenotypes in the pathogenesis of chronic wound infection^[1,2]. They go on to highlight the fundamental point that increased proliferation of exogenous microorganisms exacerbates a state believed to inhibit wound healing^[3].

It is therefore logical to focus on the importance of reducing the microbial load and preventing the introduction of new microbes during the management of chronic wounds. Historically, aseptic technique has been mandated to achieve these ends; however, the term is rarely adequately defined, nor is the practical application adequately explained. As a result, practice standards vary widely^[4-6], ultimately contributing to failures in aseptic technique and, in turn, contributing to the existence of chronic wounds.

Having a standard approach to aseptic technique is increasingly mandated internationally^[7,8], as it has been shown to improve healthcare worker knowledge and clinical practice and, in a growing number of cases, play a significant role in reducing healthcare-associated infection^[9]. One example of a robust, evidence-based clinical practice framework that has improved competency-based teaching and the

practice of aseptic technique is the now globally recognised aseptic non-touch technique initiative^[10].

Wolcott and Fletcher attempt to thoroughly address one component of the overall practice of aseptic technique: the importance of delivering effective wound irrigation. In exploring the variety of potential solutions that could be employed in this crucial activity, they touch on the controversy that surrounds sterilised saline solutions and tap water for irrigation. The most recent Cochrane review of randomised and quasi-randomised trials reports no evidence that saline and tap water are detrimental to wound healing^[11], but nor does it show that these solutions exhibit the antimicrobial effectiveness needed to help manage chronic wounds. Furthermore, there are possible water source-related problems with using tap water. For example, Gram-negative *Pseudomonas* bacteria is a well-reported issue in tap water^[12] and, given the potential for biofilm development in healthcare water systems^[13], the nature and extent of its presence in chronic wounds should be investigated further. Chronic wounds are not often swabbed, which limits an epidemiological approach to biofilm in chronic wounds. Moreover, in community settings (where the majority of chronic wounds are cared for), water quality is not routinely monitored, underappreciating the role it might play in the development of



Authors: Stephen Rowley, Clinical Director; Simon Clare, Director of Research and Development, The Association for Safe Aseptic Practice, London, UK

gram-negative-rich biofilms.

Although we cannot know the extent of delayed healing of wounds related to tap water, the potential to impede healing combined with the lack of antimicrobial activity on the parts of both tap water and saline solution mean they cannot be viewed as the most effective options for irrigation of chronic wounds. Clinicians should seek to use a solution that balances antimicrobial activity with minimal cytotoxicity as part of managing wound biofilm. Whatever wound irrigation solution is chosen and used should be in accordance with standardised aseptic technique, to decrease the risk of microorganism proliferation. A holistic approach to chronic wound care, incorporating the practical tips provided by Wolcott and Fletcher is therefore essential to working towards solving the huge problem of chronic wound prevalence. In an area short of gold-standard evidence, Wolcott and Fletcher's appraisal of both the role and application of wound irrigation will be welcomed by practitioners. WINT

References

1. Martin JM, Zenilman JM, Lazarus GS. Molecular microbiology: new dimensions for cutaneous biology and wound healing. *J Invest Dermatol* 2010; 130(1): 38–48
2. James GA, Swogger E, Wolcott R, et al. Biofilms in chronic wounds. *Wound Repair Regen* 2008; 16(1): 37–44
3. Schierle CF, De la Garza M, Mustoe TA, Galiano D. Staphylococcal biofilms impair wound healing by delaying reepithelialisation in a murine cutaneous wound model. *Wound Repair Regen* 2009; 17(3): 354–9
4. Aziz AM. Variations in aseptic technique and implications for infection control. *Br J Nurs* 2009; 18(1): 26–31
5. Flores A. Sterile versus non-sterile glove use and aseptic technique. *Nurs Stand* 2008; 23(6): 35–9
6. Rowley S, Clare S, Macqueen S, Molyneux R. ANTT v2: an updated practice framework for aseptic technique. *Br J Nurs* 2010; 19(Sup1): S5–11
7. Department of Health. *The Health and Social Care Act 2008: Code of practice on the prevention and control of infections and related guidance*, 2010. Available at: www.gov.uk/government/publications/the-health-and-social-care-act-2008-code-of-practice-on-the-prevention-and-control-of-infections-and-related-guidance
8. National Health and Medical Research Council. *Australian Guidelines for the Prevention and Control of Infection in Healthcare*, 2010. Available at: www.nhmrc.gov.au/guidelines/publications/cd33 (accessed 24.09.2014)
9. Rowley S, Clare S. Improving standards of aseptic practice through an ANTT trust-wide implementation process: a matter of prioritisation and care. *J Infect Prevent* 2009; 10(Suppl1): S18–23
10. White J. Solving the invisible problem. Poster. Presented at: Institute for Healthcare Improvement International Forum on Quality and Safety in Healthcare; Nice, France, 20–23 April 2010
11. Fernandez R, Griffiths R. Water for wound cleansing. *Cochrane Database Syst Rev* 2012; 15; 2: CD003861. doi: 10.1002/14651858.CD003861.pub3.
12. Jefferies JM, Cooper T, Yam T, Clarke SC. *Pseudomonas aeruginosa* outbreaks in the neonatal intensive care unit — a systematic review of risk factors and environmental sources. *J Med Microbiol* 2012; 61(Pt 8): 1052–61
13. Trautmann M, Lepper PM, Haller M. Ecology of *Pseudomonas aeruginosa* in the intensive care unit and the evolving role of water outlets as a reservoir of the organism. *Am J Infect Control* 2005; 33(5 Suppl 1): S41–9