>QUICK GUIDE

3M[™] Snap[™] Therapy System



Clinical Pathway to 3M™ Snap™ Therapy System

Full holistic assessment of patient and wound

Is the wound type indicated for NPWT use without contraindications¹?

Snap Therapy System is indicated for the removal of small amounts of exudate from:

- Chronic, acute, traumatic, sub-acute and dehisced wounds
- Ulcers (such as diabetic, venous or pressure)
- Surgically closed incisions
- Flaps and grafts



Wound characteristics that may be appropriate for the Snap Therapy System

- Longer-than-expected duration for the wound type
- Low-to-moderate volumes of exudate
- Thin- to medium-viscosity exudate
- Wound bed not granulating
- Shallow cavity wounds <3cm deep. For deeper wounds - consider traditional NPWT



Dressing/patient characteristics that may be appropriate for the Snap Therapy System

- Wound size <18cm x 18cm
- Wound drainage <180ml of exudate per week
- Active, ambulatory or elderly patients
- Patient wants and is able to actively engage in self-care or has a carer who can assist
- Patient has no allergy to hydrocolloid adhesives or polyurethane films



Initiate Snap Therapy System, with dressing changes a minimum of two times per week, or as exudate levels dictate. Review the dressing regimen weekly.









- Active Bleeding
- Malignand
 Untreated
- Osteomyelitis
- sels, Tendons, Nerves, Anastomosis Sites
- Active Infectio



Using your 3M[™] Snap[™] Therapy System

Prepare the wound bed and periwound skin per institutional protocol. If necessary apply a skin protectant to the surrounding skin and allow to dry.

STEP 1: Cut the foam interface to fit the size and shape of the wound. Remove backing and carefully place the $3M^{TM}$ Snap TM Advanced Dressing over the wound and seal.









STEP 2: Cut the dressing tubing to the desired length, if required. Fully insert the Tube Fitting into the tubing.





STEP 3: Connect the 3M[™] Snap[™] Cartridge to the Tube Fitting using both hands. Activate the Snap Cartridge by pushing down the Activation/Reset Key to release and remove the key from the Snap Cartridge. Cartridge may need to be primed multiple times depending on size of the dressing.

NOTE: Keep the Activation/Reset Key. DO NOT throw away.






STEP 4: Check negative pressure operation. Snap Therapy System is working properly if:


- Green capacity indicator is both visible and stationary in the chamber window
- Dressing has a "sucked down" appearance
- >> Dressing feels hard to the touch




Dressing application




 $3M^{\mathsf{TM}}\operatorname{Snap^{\mathsf{TM}}}\operatorname{Advanced}\operatorname{Dressing}\operatorname{Application}\operatorname{on}\operatorname{a}\operatorname{Heel}$



3M[™] Snap[™] Advanced Dressing Application on a Toe Amputation/Wedge



Snap Advanced Dressing Application on a Transmetatarsal Amputation (TMA)



3M[™] Snap[™] Therapy System



For the initial application, you need to order:




If required, you may also need these accessories:



Summary

The 3MTM SnapTM Therapy System is a wearable dNPWT device that combines the simplicity of advanced wound dressings with the proven efficacy of negative pressure wound therapy in a discreet design that helps preserve patient quality of life.1

- Mechanically powered and portable for patient mobility
- Delivers clinically proven -125mmHg continuous negative pressure
- Discreet and can be worn under clothing
- Silent design minimizes. sleep interruptions



NOTE: Specific indications, contraindications, warnings, precautions and safety information exist for these products and therapies.

Please consult a clinician and product instructions (as a consult a clinician and product instructions).

Please consult a clinician and product instructions for use prior to application. This material is intended for healthcare professionals.

Acceptable 2015



© Wounds International 2021 Supported by 3M www.3M.com

© 2021 3M. All rights reserved. 3M and the other marks shown are marks and/or registered marks. Unauthorized use prohibited. Used under license in Canada. PRA-PM-UK-00008 (04/21)